

LOW PRESSURE HYPERBARIC TREATMENT FOR POST SURGICAL WOUND HEALING

Loren Z. Clayman, M.D.P.A

PLASTIC AND RECONSTRUCTIVE SURGERY
DIPLOMAT-AMERICAN BOARD OF PLASTIC SURGERY

ABSTRACT

Mohs cancer surgery results in a large open wound which may be closed with a flap or skin graft or allowed to heal with secondary intention. Flaps and skin grafts often result in significant scarring of the skin. Secondary intentions allows healing in a small concentric scar. Normal healing of a defect or wound over the forehead may take up to 12 weeks to close by secondary intention. Using daily low pressure hyperbaric air chamber treatments a 3cm X 4cm defect of the forehead healed in 4 weeks. The wound edges remained soft and supple allowing fibroblastic contraction to occur significantly faster. The photographs show closure of the forehead defect at four weeks. These findings display a significant increase in healing over standard secondary intent in healing.

Mark A. Clayman, M.D.
University of Florida
Department of Surgery
Gainesville, FL

Loren Z. Clayman, M.D.
St. Vincent's Medical Center
Department of Plastic Surgery
Jacksonville, FL

Loren Z. Clayman, M.D.P.A.
St. Vincent's Medical Center
Department of Plastic Surgery
Jacksonville, FL

Mark A. Clayman, M.D.
University of Florida
Department of Surgery
Gainesville, FL

Loren Z. Clayman, M.D.
St. Vincent's Medical Center
Department of Plastic Surgery
Jacksonville, FL



Feb 1, 2000
Post Surgery



Mar 11, 2000
After daily hyperbaric treatments

Enlargement of Wound Area



Feb 4, 2000
Post Surgery - Pre hyperbarics

Protocol

Pressure: 1.2 ATA
Duration: 1 hour
Frequency: Daily for 1 month